

Public Health Reports

VOLUME 63

MAY 28, 1948

NUMBER 22

IN THIS ISSUE

Leprosy in California

Public Health Service Publications

Incidence of Communicable Disease in the U. S.



FEDERAL SECURITY AGENCY

PUBLIC HEALTH SERVICE

FEDERAL SECURITY AGENCY

Oscar R. Ewing, Administrator

PUBLIC HEALTH SERVICE

Leonard A. Scheele, Surgeon General

Division of Public Health Methods

G. St. J. Perrott, Chief of Division

CONTENTS

	Page
Leprosy in California—Danger of Infection. G. W. McCoy	705
Incidence of Communicable Disease in the United States March 28–April 24, 1948	712
Public Health Service Publications. A list of publications issued during the period July–December 1947	716

INCIDENCE OF DISEASE

United States:

Reports from States for week ended May 8, 1948, and comparison with former years	722
--	-----

Weekly reports from cities:

City reports for week ended May 1, 1948	726
---	-----

Rates, by geographic divisions, for a group of selected cities	728
--	-----

Plague infection in Arizona, New Mexico, and Washington	728
---	-----

Territories and possessions:

Panama Canal Zone—Notifiable diseases—March 1948	729
--	-----

Foreign reports:

Canada—Provinces—Communicable diseases—Week ended April 17, 1948	730
--	-----

Northern Rhodesia—Smallpox	730
----------------------------------	-----

World distribution of cholera, plague, smallpox, typhus fever, and yellow fever—

Cholera	731
---------------	-----

Plague	732
--------------	-----

Smallpox	732
----------------	-----

Typhus fever	734
--------------------	-----

Yellow fever	735
--------------------	-----

Deaths during week ended May 1, 1948	735
--	-----

Public Health Reports

Vol. 63 • MAY 28, 1948 • No. 22

Printed With the Approval of the Bureau of the Budget as Required by Rule 42
of the Joint Committee on Printing

LEPROSY IN CALIFORNIA—DANGER OF INFECTION

G. W. McCoy, M. D.

Reports on the prevalence of leprosy often do not disclose where the infection was acquired. This is important in connection with control measures if the case is found in an area in which transmission to the public is likely to occur. With new interest being taken in this disease by health authorities in the United States, it is desirable to know the danger of spread in different areas. This seemed especially important for California where many cases have been reported. Among 475 cases reported up to 1940, probably not more than 14 had been infected in the State (1). The present investigation was undertaken to obtain more comprehensive and more recent data. California stands almost alone among the States in being geographically situated for receiving infection from two general sources. The proximity to Mexico, which has many cases, has resulted in the entrance of a very considerable number of cases from that country. The State is the most convenient port of entry for Hawaii, Japan, and other Pacific islands, as well as China, which has resulted in the importation of a smaller but considerable number of cases. These undoubtedly included active cases and probably a larger number in the incubation stage.

At the time California became a part of the United States (1850), the natives, excluding Indians, were designated as Hispano-Californians (Mexicans) and constituted a large part of the population. A number of native Hawaiians came to the State soon after discovery of gold (1848), and at about the same time Chinese began to come in large numbers. Doubtlessly, both immigrations brought leprosy with them. The first readily available reference to the disease in California is a report by the jail physician of San Francisco in 1877, who mentioned treatment of 3 cases of "leprosis." A little later the State health officer of the day, Dr. H. S. Orme, in an annual report (2) included the following statistical data on lepers in San Francisco from 1871 to 1890, based on California State Board of Health reports from 1886 to 1894:

Year of admission:	Cases	Year of admission:	Cases	Year of admission:	Cases
1871.....	1	1878.....	13	1885.....	7
1872.....	1	1879.....	14	1886.....	6
1873.....	1	1880.....	10	1887.....	3
1874.....	6	1881.....	2	1888.....	3
1875.....	9	1882.....	12	1889.....	12
1876.....	3	1883.....	11	1890.....	5
1877.....	0	1884.....	9	Total.....	¹ 128

¹ Of this total, 115 were Mongolians, 12 whites and 1 mixed; 120 were males and 8 females.

Birthplaces were recorded as follows:

China.....	114	England.....	1	Japan.....	1
Honolulu.....	1	Sweden.....	1	Germany.....	1
United States.....	3	France.....	1	Mexico.....	1

These figures show an overwhelming predominance of Chinese and an extraordinary excess of males. Nearly everywhere the number of male victims is much larger than females, the ratio generally being about 2:1. The very great excess shown in California in those early years doubtlessly resulted from confining Chinese immigration largely to males. Dr. Orme also wrote "Long ago the people of California recognized the danger of planting leprosy on this coast through Chinese immigration, and for more than 15 years legislation gave abundant authority for its exclusion and repression. Section 2952 of the Political Code reads: 'It shall not be lawful for lepers or persons affected with leprosy, or elephantiasis, to live in ordinary intercourse with the population of this State; but all persons shall be compelled to inhabit such lazarettos, or leper's quarters, as may be assigned to them by the Board of Supervisors of the city or county in which they shall be domiciled or settled; and the Boards of Supervisors are vested with power and are required to make all necessary provisions for the separation, detention, and care of lepers or persons affected with leprosy, or elephantiasis, settled or domiciled in their respective cities or counties.'

"In 1883," Dr. Orme continued, "the Board of Supervisors of San Francisco supplemented the above act by an order which forbids positively the landing of lepers from any ship, their transfer to another vessel, and their harboring by any person outside the lazaretto." There is nothing to indicate that the cases referred to by Dr. Orme gave rise to any new infections.

It has long been known that California furnishes a large proportion of the cases of leprosy reported in the United States and, according to Hopkins and Faget (3), the State with 207 cases ranks third in the number of cases sent to the National Leprosarium at Carville, La., being exceeded only by Louisiana with 596 cases and Texas with 214 for the period 1921-44. Until this study, there seems to have been no special attempt to determine how many, if any, of the cases re-

ported for California were infected there. Comparatively recently, when most health authorities began the adoption of new and more intelligently directed policies in dealing with this disease, it became important to know where leprosy is communicated from the sick to the well, in addition to where it is found. Florida, Louisiana, and Texas have long been recognized as areas in which leprosy spreads enough to be a public health problem and in recent years there has been a tendency to group California with these States as an "endemic" area. With the object only of ascertaining the status of the State as an area in which this disease was transmitted, this study was made. When data on any case indicated that in all probability the infection had been acquired elsewhere than in California, no further attention was given to it. This course was adopted because there was no intention of making a complete study, but only of the cases that could reasonably be regarded as having been infected in this State.

When this investigation began in California in July 1947, the State Health Department was engaged in a general statistical study of leprosy in California which was to be prepared for publication. The data for that study were generously placed at my disposal and found of much value. The figures were for the period 1906-1947, and covered a total of approximately 500 cases reported in the State.

The tracing of source and place of infection in communicable diseases usually is carried out by ascertaining the place and the time of exposure, giving consideration to the incubation period. Perhaps the best illustration of this is to be found in tracing the source of venereal disease in a control program. Another example is smallpox. Look for the infecting smallpox patient among the victim's contacts 12 to 14 days previous to onset. If his contacts can be traced, the infecting patient should be found.

It is the widely accepted view that leprosy is usually acquired in the early years of life. Determining the source and the place of infection in leprosy is, therefore, often difficult and frequently impossible, chiefly for two reasons: First, the manifestations in the infecting patient may be obscure and the diagnosis in the new patient uncertain for long periods, even many years; secondly, the incubation period is long and varying and may average from 8 to 10 years. However, some authorities feel that it may be prolonged to 25 years or longer. In this study, if the patient had lived in Hawaii, or the Philippines, or a similar recognized area of high leprosy prevalence, this was regarded as the place where the disease probably was acquired. A few years' residence in either of these Pacific Island groups was considered sufficient to attribute the infection to that area. If the patient had been born in Mexico, and had spent the early years of life in that country, Mexico was considered the probable place of infection even if a longer period later in life had been spent in California.

The following procedures have been utilized in this study. Patients admitted from California to the National Leprosarium at Carville, La., and who remained there, were interviewed in May 1947. Among 74 such cases 7 could be assigned to California as the place of infection. The records of the California State Department of Health were studied in July and August 1947. They were so complete that not much difficulty was encountered in allocating cases to a place of probable infection. The State records included most of the patients who had been interviewed at Carville in May 1947.

In recent years, the attitude of California health authorities in general has been very enlightened and progressive with respect to the public health management of the disease. While patients regarded as possible (or probable) sources of infection have been sent to the National Leprosarium at Carville, La., very little compulsion has been employed. Patients judged not to be a menace to those about them often have been allowed to remain at home, in local hospitals, or under the care of local physicians. Patients of Mexican origin who preferred to return to their native land were permitted to and informal arrangements for receiving the deportees were even facilitated by the State or local authorities with Mexican authorities.

CASE STUDIES

The family groups with leprosy (excluding marital) appeared to present the clearest evidence of infection within the State. Three of these are of special interest as the cases charged to the State never had been outside California:

(1) K family: A mother, born in Japan, came to the United States when 14 years old and developed leprosy 15 years later. At the time of diagnosis she was classified as an active case of leprosy. Her husband, also born in Japan and the father of her 6 sons, was rated as an old inactive case, but the evidence in his case is not conclusive. The sons who contracted the disease were their third, age 12, the fifth, age 9, and the sixth, age 8.

(2) S family: The first case in this family was the father who died of leprosy in 1928 and never had been out of the State. Ten years later a son, age 17, was infected. He also had never been out of California.

(3) F family: The father in this family had military service in the Philippine Islands (1899-1900) and had developed leprosy after his return, dying in 1912. His two sons, one born in 1901 and the other in 1902, developed the disease, one at age 16 and the other at 17. Both had lived at home with the father until 18 months before his death. A daughter is said to have escaped infection.

Another familial case centers about Mrs. M. P., age 40, who was found to have advanced leprosy in 1932. She was born and probably infected in Mexico. She died of pneumonia apparently while hospitalized for leprosy. A short time later a daughter, I. P., age 6, who had never lived out of California, was discovered to have early leprosy.

ADULT INFECTIONS

Almost as clear examples of adult infection within the State are at least two patients who had lived only in California, except for limited periods in areas where the disease never has been known to be transmitted:

A young adult male (G. P.) who had been out of California only for 3 years while serving as a sailor in Alaska and Alaskan waters; a woman (M. B.), age 48, whose only residence outside California was 7 years spent in Utah. Neither Alaska nor Utah is considered as a probable area of leprosy transmission.

Marital infections.—In the three instances in which husband and wife were reported to have leprosy, the marital partner first to present evidence of the disease was regarded, perhaps somewhat arbitrarily, as the source of infection for the mate. In one family it was not possible to reach any reasonably clear conclusion on this point. In all of the marital infections, the evidence is not so convincing since both partners had been born outside California. If the view of infection of one from the other had not been adopted, it would have been necessary to assume the coincidence that both partners had been infected abroad. Three marital infections in so small a total is a much larger number than is to be expected and throws doubt on some, or all of them.

The K. family: both partners previously mentioned were born in Japan and spent their early years there. It was impossible to be certain that either one was infected in California, but the probability appears to point in that direction. The alternative is to consider that both were infected in Japan.

The S. family: Mrs. M. S. was born in Mexico and came to California at the age of 28. Nine years after coming to the State she was found to have well-advanced leprosy of the lepromatous type. Her husband, S. S., age 49, was born in Mexico. He spent the first 5 years of his life in Mexico and the remaining 44 years in California. The disease appeared about 1 year before the diagnosis of the maculo-anesthetic form was made; that is, 43 years after coming to the United States.

The R. family: Mr. J. R., age 49, was born in Mexico. He came to California at the age of 31 and was reported as having leprosy 18 years later. The disease was classified as "mixed" of long duration. His wife Jo. R., age 41, was born in Mexico and had lived in California 14 years. Her case was recorded as "macular leprosy, early."

Certain data on the 23 cases important to the present inquiry are shown in the following tabulation:

TABLE 1.—*Leprosy infection in Californians who never left the State, probably acquired in California*

Year reported	Initials	Sex	Location	Source of infection
1919	T. F.	Male	Stockton	Father.
1919	W. F.	do.	do.	Do.
1925	H. S.	do.	Walnut Grove	Unknown.
1932	I. P.	Female	Tulare County	Mother.
1938	J. S.	Male	Walnut Grove	Father.
1941	A. K.	do.	Sacramento	Mother.
1941	W. K.	do.	do.	Do.
1941	K. K.	do.	do.	Do.

TABLE 2.—*Leprosy infection in persons never out of State except to areas where disease is not known to be communicated*

Year	Initials	Sex	Location	Birthplace	Residence	Number of years outside State
1921	G. P.	Male	San Leandro	California	Alaska	3
1943	M. B.	Female	Angels Camp	California	Utah	7

TABLE 3.—*Cases where marital partner is probable source of infection*

Year reported	Initials	Sex	Location	Birthplace	Probable source of infection
1929	Y. K.	Female	Sacramento	Japan	Husband.
1939	S. S.	Male	Carlsbad	Mexico	Wife.
1941	J. R.	Female	Mount Shasta	do.	Husband.

TABLE 4.—*Cases believed to indicate infection in the State*

Year	Initials	Sex	Location	Birthplace	Residence outside the State
1926	M. L.	Female	Decoto	(¹)	(¹)
1930	E. L.	Male	Los Angeles	Calif.	None.
1931	J. S.	do.	Los Angeles area.	do.	Mexico, 1 day (7 years prior to onset).
1935	C. R.	do.	Hanford	do.	None.
1935	R. S.	do.	San Francisco	do.	(¹)
1939	A. M.	do.	do.	do.	Pennsylvania, Oregon, Montana, Vancouver 5 years.
1943	A. S.	do.	(¹)	do.	Mexico before age 9.
1945	D. M.	do.	Oakland	Spain	Spain first year of life.
1946	R. M.	Female	Fresno	Arizona	Arizona first year of life.
1947	E. H.	do.	Los Angeles area.	Ohio	Ohio for first 50 years.

¹ Not known.

In comparing the incidence of intrastate infected cases with the total number that were reported for periods for which data are available, the figures are as follows:

Period	Total cases reported	Probably infected in State
1913-1916.....	43	0
1917-1921.....	112	3
1922-1926.....	100	2
1927-1931.....	102	3
1932-1936.....	68	3
1937-1941.....	54	7
1942-1946.....	49	4

The period 1937-1941 includes the three children in one family. One case of within-State infection was omitted from this table as it was reported in 1947.

The figures show a tendency of total reported cases to be falling in recent years. The largest number for any one year was 28 in 1921. Intrastate infections do not vary much for the period under consideration.

CLASSIFICATIONS OF AREAS OF INFECTIVITY

The prevalence of leprosy in the United States by classifying areas is as follows:

(1) *Highly endemic*.—All or the great majority of cases being infected in the State—Louisiana and Florida.

(2) *Markedly endemic*.—A large proportion of the cases being infected in the State—Texas.

(3) *Mildly endemic*.—A very small proportion of all cases being infected in the State—Minnesota.

(4) *Feebly endemic*.—Only occasional cases occurred at long intervals—South Carolina.

In such a grouping California would fall in the class of mildly endemic along with Minnesota. Minnesota has had 7 cases of local origin among a total of 100 or more, while in California the figures are approximately 23 among about 500—a rather suggestive similarity between the two States.

When the data are examined from the point of view of locality in the State where infection occurred, it is found that eight of the cases came from a comparatively small area near the middle of the State as follows:

Sacramento—4 cases in one family—mother, 3 children

Walnut Grove—2 cases in one family—father, 1 child

Stockton—2 cases in one family—2 brothers

These are from an area with a radius of about 30 miles. Of the 8 cases, 6 were children in families with a leprous parent.

The experience of California seems to agree to a considerable extent with that of Minnesota where the number of cases acquiring the

disease in the State was far outnumbered by imported cases. There is one important difference between the two States. In California, leprosy continues to be introduced with the importation of patients or persons in the incubation period from Mexico and the Pacific area, while in Minnesota, an end to the admission of new cases came about the beginning of the 20th century.

SUMMARY

An investigation having a very limited objective, shows the following: That 23 persons in the present century have acquired leprosy in California. Of these, seven never had been out of the State. About these seven there can be no doubt as to the place of infection. The remainder are less positively attributed to infection in the State but this is believed to be established beyond reasonable doubt.

The majority of all cases are due to infection in Mexico, China, and the Pacific Islands.

California is to be regarded as an area in which the likelihood of transmission of leprosy is small, except for children born of parents, one or both of whom have the disease.

ACKNOWLEDGMENT

The author wishes to express his indebtedness for aid to Miss Ida M. Stevens and Mr. Guy P. Jones, of the California State Health Department; to R. C. Williams, Assistant Surgeon General; to F. A. Johansen, Medical Director; and M. V. Hargett, Senior Surgeon, Public Health Service.

REFERENCES

- (1) McCoy, G. W.: Observations on the epidemiology of leprosy, *Pub. Health Rep.* **57**: 1935 (1942).
- (2) Orme, H. S.: Report of State Board of Health of California June 30, 1888-June 30, 1890, page 211.
- (3) Hopkins, Ralph, and Faget, G. H.: Recent trends in leprosy in the United States, *J. A. M. A.* **126**: 937 (1944).

INCIDENCE OF COMMUNICABLE DISEASES IN THE UNITED STATES

March 28-April 24, 1948

The accompanying table summarizes the incidence of nine important communicable diseases, based on weekly telegraphic reports from State health departments. The reports from each State for each week are published in *PUBLIC HEALTH REPORTS* under the section "Incidence of Disease." The table gives the number of cases of these diseases for the 4 weeks ended April 24, 1948, the number reported for the corresponding period in 1947, and the median number for the years 1943-47.

DISEASES ABOVE MEDIAN INCIDENCE

Influenza.—The reported number of cases of influenza dropped from 25,459 during the preceding 4 weeks to 10,095 for the 4 weeks ended April 24. Of the total cases Texas reported 4,393, South Carolina 1,486, and Virginia 1,138—70 percent of the total cases occurred in those 3 States. The recent rise of this disease has been confined to States in the Southern and Western sections of the country. In the East South Central, Mountain and Pacific sections the incidence has dropped nearly to the level of the preceding 5-year median, but in the South Atlantic and West South Central sections the numbers of cases are still relatively high. The incidence for the country as a whole was less than 10 percent of the 1947 figure for the same weeks, but it was about 17 percent above the 1943-47 median incidence. The peak of the recent outbreak was reached during the month of January while the peak of the 1947 epidemic was not reached until March, with very little drop in the number of cases during the month of April.

Poliomyelitis.—While the number of cases of poliomyelitis was not large, it represented an increase of more than 10 percent over the median for the preceding 5 years, and with the exception of the year 1945, when 128 cases were reported for these same 4 weeks, the current incidence was the highest for the corresponding weeks in the 20 years for which data are available in this form. Each section of the country except the North Atlantic, East South Central, and Pacific sections reported an excess of cases over the normal seasonal median. While an increase of this disease may be expected at this season of the year, in most preceding years the increase has not occurred until sometime during the month of May; the lowest incidence of the season has normally been reached during the 4 weeks corresponding to the period under consideration.

DISEASES BELOW MEDIAN INCIDENCE

Diphtheria.—For the 4 weeks ended April 24 there were 610 cases of diphtheria reported. The median number for the preceding 5 years was 922. For the country as a whole the current incidence was the lowest on record for this period. In the Mountain section the number of cases (75) was 50 percent above the seasonal expectancy, but in all other sections the incidence was relatively low.

Measles.—The number of reported cases of measles rose from 83,160 during the preceding 4 weeks to 102,680 during the current 4-week period. The incidence was 3.6 times that reported for the corresponding period in 1947, but it was slightly below the median for the preceding 5 years (104,809 cases). Significant increases over the normal seasonal incidence were reported from the East North Central,

West South Central, Mountain, and Pacific sections. In other sections the incidence was about the same or below the preceding 5-year medians.

Meningococcus meningitis.—The number of cases (285) of meningococcus meningitis was less than 75 percent of the number reported for the corresponding period in 1947 and about 36 percent of the median for the preceding 5 years. In the Mountain section the incidence was the same as the median but in all other sections the incidence was considerably below the 1943-47 median. For the country as a whole the current incidence was the lowest since 1941 when 225 cases were reported for these same weeks. The current incidence compares very favorably with the average number of cases (approximately 275) for non-epidemic years, the 1943-47 median contains 3 years of unusually high meningitis incidence.

Scarlet fever.—This disease continued on its downward trend, the number of cases (8,312) reported for the 4 weeks ended April 24 being below the record low incidence of 1947 (9,898 cases). In each section of the country the current incidence was considerably below the normal seasonal incidence. This disease has been on a steady decline since 1944 and for this particular period the incidence was the lowest in the 20 years for which these data are available.

Smallpox.—Kansas and North Carolina each reported 2 cases of smallpox and Nebraska, Wyoming, and Arizona one each during the current 4 weeks. There were 38 cases reported during the corresponding period in 1947 and the median for the preceding 5 years was 54 cases. During this period in 1947 there was a slight outbreak of smallpox around New York City, the first outbreak in New York State since 1939, and 13 cases were also reported from Texas during these weeks. For the country as a whole the current incidence is the lowest on record for this period.

Typhoid and paratyphoid fever.—The number of cases of these diseases increased from 166 cases during the preceding 4 weeks to 225 during the current 4-week period. The number of cases was 1.4 times the 1947 incidence for the same weeks and slightly below the median for the preceding 5 years. The greatest increases over the 1947 figures occurred in the South Atlantic and West South Central sections with minor increases in all other sections except the New England, Mountain, and Pacific sections. Compared with the preceding 5-year median, the current incidence was higher than the median in the New England, West North Central, South Atlantic, and West South Central sections; in other sections the numbers of cases were about the same or below the medians.

Whooping cough.—For the 4 weeks ended April 24 there were 8,001 cases of whooping cough reported, as compared with 10,545 in 1947 and a median of 10,035 for the corresponding weeks in the preceding

5 years. In the West South Central section the number of cases (2,172) was 1.9 times the seasonal expectancy and a few more cases than might normally be expected occurred in the West North Central and Mountain sections, but in all other sections the incidence was relatively low.

MORTALITY, ALL CAUSES

For the 4 weeks ended April 24 there were 37,407 deaths from all causes reported to the National Office of Vital Statistics by 93 large cities. The median number for the years 1945-47 was 36,731 deaths. During the first 2 weeks of the 4-week period the number of deaths was considerably above the preceding 3-year median, but during each of the last 2 weeks the number was lower than the median. For the entire 4-week period the increase over the median was about 1.8 percent.

Number of reported cases of 9 communicable diseases in the United States during the 4-week period March 28-April 24, 1948, the number for the corresponding period in 1947, and the median number of cases reported for the corresponding period, 1945-47

Division	Current period	1947	5-year median	Current period	1947	5-year median	Current period	1947	5-year median
	Diphtheria			Influenza ¹			Measles		
United States.....	610	922	922	10,095	120,721	8,650	102,680	28,280	104,809
New England.....	21	47	32	7	225	108	5,840	7,076	7,487
Middle Atlantic.....	63	165	163	72	119	72	20,492	4,328	20,955
East North Central.....	86	118	166	80	3,868	427	29,616	5,055	26,395
West North Central.....	39	85	70	114	14,564	108	7,401	1,444	7,441
South Atlantic.....	124	170	131	2,878	36,811	2,486	5,349	4,369	7,035
East South Central.....	54	88	85	635	9,740	606	2,161	1,465	3,182
West South Central.....	90	127	152	5,279	48,582	4,277	11,332	2,070	5,963
Mountain.....	75	49	50	664	4,946	609	5,269	1,401	4,643
Pacific.....	58	73	125	366	1,866	344	15,220	1,072	7,945
	Meningococcus meningitis			Poliomyelitis			Scarlet fever		
United States.....	285	383	794	126	112	111	8,312	9,898	17,096
New England.....	10	12	39	0	2	2	916	899	2,211
Middle Atlantic.....	52	57	155	16	20	19	2,374	2,769	5,679
East North Central.....	61	85	152	19	13	10	2,645	2,971	4,247
West North Central.....	21	37	72	19	9	5	595	824	1,552
South Atlantic.....	48	68	122	16	9	9	463	645	1,340
East South Central.....	37	38	68	6	9	9	187	419	475
West South Central.....	25	45	73	31	18	20	197	195	492
Mountain.....	10	6	10	5	1	3	287	395	855
Pacific.....	21	35	103	14	31	21	648	781	1,061
	Smallpox			Typhoid and paratyphoid fever			Whooping cough		
United States.....	7	38	54	225	161	241	8,001	10,545	10,035
New England.....	0	0	0	24	24	9	532	824	932
Middle Atlantic.....	0	13	0	16	15	37	915	1,738	1,738
East North Central.....	0	3	7	24	18	27	1,199	2,110	1,476
West North Central.....	3	2	8	15	13	10	516	318	318
South Atlantic.....	2	0	2	58	26	52	1,084	1,363	1,533
East South Central.....	0	5	4	22	16	23	326	588	463
West South Central.....	0	15	8	48	25	40	2,172	2,313	1,160
Mountain.....	2	0	2	1	4	15	659	324	474
Pacific.....	0	0	6	17	20	20	508	967	967

¹ New York, North Carolina, and Pennsylvania excluded; New York City and Philadelphia included.

PUBLIC HEALTH SERVICE PUBLICATIONS**List of Publications Issued During July-December 1947**

The purpose of this list is to provide a complete and continuing record of Public Health Service publications for reference use by librarians, scientific workers, and others interested in particular fields of public health work, and not to offer the publications for indiscriminate free distribution.

Single sample copies are available from the Public Inquiries Section, Office of Health Information, Public Health Service, Washington 25, D. C.

Quantities may be obtained from the Superintendent of Documents, Government Printing Office, Washington 25, D. C., at prices shown, with a reduction of 25 percent on lots of 100 copies or more of a single publication.

The publications marked with an asterisk (*) may be obtained only by purchase.

Periodicals

- *Public Health Reports (weekly), July-December, vol. 62, Nos. 27 to 52. Pages 969 to 1812. 10 cents a number. Subscription price \$4 a year.
- Extracts from Public Health Reports (monthly), July-December, Tuberculosis Control Issues Nos. 17 to 22. 30 pages each. 10 cents a number. Subscription price \$1 a year.
- *The Journal of Venereal Disease Information (monthly), July-December, vol. 28, Nos. 7 to 12. Pages 129 to 298. 10 cents a number. Subscription price 75 cents a year.
- *Journal of the National Cancer Institute (bimonthly), August-December, vol. 8, Nos. 1 to 3. Pages 1 to 159. 40 cents a number. Subscription price \$2 a year.
- Public Health Engineering Abstracts (monthly), July-December, vol. XXVII, Nos. 7 to 12. 32 pages each. No sales stock.
- Industrial Hygiene Newsletter (monthly), July-December, vol. 7, Nos. 7 to 12. 16 pages each. 10 cents a number. Subscription price \$1 a year.
- National Negro Health News (quarterly), July-December, vol. 15, Nos. 3 and 4. 24 pages each. No sales stock.

Reprints from the Public Health Reports

- 2795. "Albumin-bacterioplasm conjugates" with special reference to the etiology of rheumatic fever. By Mark P. Schultz and Edythe J. Rose. July 11, 1947. 14 pages. 5 cents.
- 2796. Public Health Service publications. A list of publications issued during the period July-December 1946. July 11, 1947. 5 pages. 5 cents.
- 2797. Comparison of the spirocheticidal activity of arsenphenamines and phenarsines (Arsenoxides) in experimental syphilis. By T. F. Probey. July 18, 1947. 8 pages. 10 cents.
- 2798. Toxic effects of tetranitromethane, a contaminant in crude TNT. By Rudolph F. Sievers, Edward Rushing, Helen Gay, and A. R. Monaco. July 18, 1947. 13 pages. 5 cents.
- 2799. Nutrition studies. I. Description of physical signs possibly related to nutritional status. By Harold R. Sandstead and Richmond K. Anderson. July 25, 1947. 13 pages. 5 cents.

2800. Mental hygiene in public health. By Paul V. Lemkau. August 8, 1947. 12 pages. 5 cents.
2801. Preliminary report on some organic materials as tick repellents and toxic agents. By James M. Brennan. August 8, 1947. 4 pages. 5 cents.
2802. State planning for participation in the National Mental Health Act. By R. H. Felix. August 15, 1947. 9 pages. 5 cents.
2803. DDT in oil as a mosquito larvicide. By Henry A. Johnson and Willis L. Goodman. August 15, 1947. 10 pages. 5 cents.
2804. Loss of virulence of *Treponema pallidum* during processing of dried blood serum. By T. F. Probey. August 15, 1947. 4 pages. 5 cents.
2805. Control of rabies. Report by the Committee on Public Health Relations of the New York Academy of Medicine. August 22, 1947. 24 pages. 10 cents.
2806. The family and dental disease. By Henry Klein. August 29, 1947. 8 pages. 5 cents.
2807. Effects of DDT mosquito larviciding on wildlife. Part II. Effects of routine airplane larviciding on bird and mammal populations. By Arnold B. Erickson. August 29, 1947. 11 pages. 5 cents.
2808. Dental services received by children in a New York clinic. By Isidore Altman. September 19, 1947. 18 pages. 10 cents.
2809. Studies on the air transmission of micro-organisms derived from the respiratory tract. By H. du Buy, F. A. Arnold, and B. J. Olson. September 26, 1947. 24 pages. 10 cents.
2810. A case of Q fever probably contracted by exposure to ticks in nature. By Carl M. Eklund, R. R. Parker, and David B. Lackman. September 26, 1947. 4 pages. 5 cents.
2811. Isolation of poliomyelitis virus from human serum by direct inoculation into a laboratory mouse. By Hilary Koprowski, Thomas W. Norton and Walsh McDermott. October 10, 1947. 10 pages. 5 cents.
2812. The effect of topically applied fluorides on dental caries experience. By Donald J. Galagan and John W. Knutson. October 10, 1947. 8 pages. 5 cents.
2813. Recovery of ornithosis virus from pigeons in Baltimore, Md. By Dorland J. Davis and C. Leroy Ewing. October 10, 1947. 5 pages. 5 cents.
2814. The effect of morphine addiction on blood, plasma, and "extra-cellular" fluid volumes in man. By Harris Isbell. October 17, 1947. 16 pages. 5 cents.
2815. Electron microscopy of tooth structure by the shadowed collodion replica method. By David B. Scott and Ralph W. G. Wyckoff. October 17, 1947. 8 pages; 8 illustrations. 10 cents.
2816. Statistical activities in State health departments. By Daniel D. Swinney. October 24, 1947. 12 pages. 5 cents.
2817. Sickness absenteeism among male and female industrial workers 1937-46, inclusive. By W. M. Gafafer. October 24, 1947. 4 pages. 5 cents.
2818. Vertebral body trephine. A preliminary report. By A. A. Michele and F. J. Krueger. August 8, 1947. 4 pages; 1 illustration. 5 cents.
2819. Births, infant mortality and maternal mortality in the United States—1944. By Nancy J. Brombacher. October 31, 1947. 20 pages. 10 cents.
2820. The role of health education in a public health program. By Mayhew Derryberry. November 14, 1947. 13 pages. 10 cents.
2821. Naturally occurring histoplasmosis in *Mus musculus* and *Rattus norvegicus*. By Chester W. Emmons, Joseph A. Bell, and Byron J. Olson. November 14, 1947. 5 pages. 5 cents.

2822. Opportunities for public health in disability insurance programs. By Milton I. Roemer. November 21, 1947. 12 pages. 5 cents.
2823. Relationship between per capita income and mortality, in the cities of 100,000 or more population. By Marion E. Altenderfer. November 28, 1947. 12 pages. 5 cents.
2824. Contributions of the UNRRA sanitary engineering program to international health. By Frederick F. Aldridge. December 12, 1947. 12 pages. 5 cents.
2825. A report on the histopathology of the cutaneous lesions of a case of rickettsialpox. By Lloyd R. Hershberger and Robert J. Huebner. December 12, 1947. 5 pages; 3 illustrations. 5 cents.
2826. Dental caries prevalence and tooth mortality. A study of 24,092 Georgia children in 12 communities. By Thomas L. Hagan. December 19, 1947. 16 pages. 10 cents.
2827. Pteroylglutamic acid ("folie acid"), liver extract, and amino acids in the treatment of granulocytopenia in rats. By Floyd S. Daft. December 26, 1947. 8 pages. 5 cents.
2828. A serum protection test in tularemic infections in white rats. By Carl L. Larson. December 26, 1947. 8 pages. 5 cents.

Supplements to Public Health Reports

196. An evaluation of neurologic symptoms and findings occurring among TNT workers. By Rudolph F. Sievers. 1947. 26 pages. 20 cents.
197. Thallium. A review and summary of medical literature. By Francis F. Heyroth. 1947. 23 pages. 10 cents.
198. The incidence of rheumatic fever as recorded in general morbidity surveys of families. By Selwyn D. Collins. 1947. 59 pages. 15 cents.
199. Food handlers' schools in Hawaii. By B. J. McMorrow and F. A. Schramm. 1947. 37 pages. 17 illustrations. 15 cents.
200. Relation of plants to malaria control in Puerto Rico. By Harry D. Pratt. 1947. 39 pages; 10 illustrations. 10 cents.
201. Laboratory control of water supplies. By H. W. Streeter. 1947. 15 pages. 10 cents.
202. Problems of interpretation of the data of rodent-ectoparasite surveys and studies of rodent ectoparasites in Honolulu, T. H., Savannah, Ga., and Dothan, Ala. By LaMont C. Cole and Jean A. Koepke. 1947. 71 pages. 20 cents.
203. The toxicity and potential dangers of zinc phosphide and of hydrogen phosphide (phosphine). (A review of the nonsecret, nonconfidential and nonrestricted literature.) By W. F. von Oettingen. 1947. 17 pages. 10 cents.
180. Directory of State and Territorial health authorities, 1947. (1947 revision.) 51 pages. 15 cents.

Public Health Bulletins

296. Manual of recommended water-sanitation practice recommended by the United States Public Health Service 1946. 1947. 40 pages. 15 cents.
297. A review of the literature relating to affections of the respiratory tract in individuals exposed to cotton dust. By B. H. Caminita, William F. Baum, Paul A. Neal, and R. Schneider. 1947. 86 pages. 25 cents.
298. Health of arc welders in steel ship construction. A survey made in cooperation with United States Maritime Commission and United States Navy. By Waldemar C. Dreessen, Hugh P. Brinton, Robert G. Keenan, Thalbert R. Thomas, Edwin H. Place, and James E. Fuller. 1947. 200 pages; 16 illustrations. 55 cents.

National Institute of Health Bulletin

189. Studies on schistosomiasis. By Willard H. Wright, Eloise B. Cram, Elmer G. Berry, Paul A. Ward, Dorothy Travis, Ruth E. Rue, Virginia S. Files, Myrna F. Jones, William B. Figgat, Frederick J. Brady, Walter L. Newton, S. R. Weibel, Harold B. Warren, Mary Louise Steinle, Mirriel S. Hummel, M. O. Nolan, Elizabeth Rogers Mann, Helen M. Churchill, John Bozicevich and Helen M. Hoyem. 1947. 212 pages. 50 cents.

Miscellaneous Publication

36. Better health for your community. Revised 1947. 12 pages. 5 cents.

Health Education Series

15. Menopause. 1947. 3 pages. 5 cents; \$1 per 100.
 20. Rocky Mountain spotted fever. 1947. 4 pages. 5 cents; \$1 per 100.
 24. Measles. 1947. 4 pages. 5 cents; \$1 per 100.
 27. Smallpox. 1947. 2 pages. 5 cents; \$1 per 100.
 28. Bronchial pneumonia. 1947. 2 pages. 5 cents; 75 cents per 100.

Community Health Series

- From hand to mouth, No. 3. Revised 1947. 48 pages. 15 cents.

Unnumbered Publications

- Index to Public Health Reports, vol. 62. Part 1, January-June 1947. 15 pages. 5 cents.
 The physician in the U. S. Public Health Service. 1947. 20 pages, illustrated. 15 cents.
 Free medical care for merchant seamen. 1947. 4 pages, illustrated. 5 cents.
 Compilation of Public Health Service Regulations. Reprinted from Federal Register. September 16, 1947. 74 pages. (For official use only.)
 Compilation of Public Health Service Regulations. Supplements Nos. 1-5, September 17 through October 11, 1947. 6 pages. (For official use only.)
 Compilation of Public Health Service Regulations. Supplement No. 6, October 22, 1947. 27 pages. (For official use only.)
 Compilation of Public Health Service Regulations. Supplements Nos. 7-10, October 24 through November 22, 1947. 2 pages. (For official use only.)

Reprints from the Journal of the National Cancer Institute

59. Inhibition of cathepsins of normal calf spleen and thymus. By Mary E. Maver and Antoinette Greco. October 1947. 4 pages. No sales stock.
 60. Subcutaneous sarcomas in mice implanted with hydrocarbon-cholesterol pellets. By Michael B. Shimkin and Rose S. Wyman. October 1947. 4 pages. No sales stock.
 61. Vascular reactions of normal and malignant tissues in vivo. II. The vascular reactions of the normal and neoplastic tissues of mice to a bacterial polysaccharide from *Serratia marcescens* (*Bacillus prodigiosus*) culture filtrates. By Glenn H. Algire, Frances Y. Legallais and Helen D. Park. October 1947. 10 pages; 4 illustrations. No sales stock.
 62. Pulmonary-tumor induction by transplacental exposure to urethane. By C. D. Larsen. October 1947. 8 pages. No sales stock.
 63. Enzymatic hydrolysis of homologous amino acid amides in normal and neoplastic tissues. By Maurice Errera and Jesse P. Greenstein. October 1947. 5 pages. No sales stock.
 64. Note on benzoylarginineamidase activity in extracts of rat liver and hepatoma. By Jesse P. Greenstein and Florence M. Leuthardt. October 1947. 2 pages. No sales stock.

65. Effects of methylcholanthrene in pregnant mice. By L. C. Strong and W. F. Hollander. October 1947. 4 pages. No sales stock.
66. Procedure for the fixation, staining, and mounting of whole mounts from tissue cultures grown in Carrel D 3.5 flasks. By Wilton R. Earle. October 1947. 7 pages. No sales stock.

Reprints from The Journal of Venereal Disease Information

284. Some significant aspects of venereal-disease research. By J. F. Mahoney. July 1947. 4 pages. 5 cents.
285. The quantitative Kahn test. By Ad Harris. July 1947. 4 pages. 5 cents.
286. A revised note on quantitative Kahn tests employing 0.9 and 2.5 percent salt-solution systems. By Reuben L. Kahn. July 1947. 2 pages. 5 cents.
287. Preservation of sheep red cells for complement-fixation tests. I. An improved method. By J. Portnoy, H. N. Bossak, and Ad Harris. July 1947. 4 pages. 5 cents.
288. Treatment of the syphilitic pregnant woman with penicillin in oil-beeswax: A comparison with results obtained using aqueous sodium penicillin. By Norman R. Ingraham, Elizabeth Kirk Rose, Herman Beerman, Virgene S. Wammock, John H. Stokes, and Paul Gyorgy. August 1947. 8 pages. 5 cents.
289. Socioeconomic aspects of granuloma inguinale. By Robert B. Greenblatt. Granuloma inguinale: Streptomycin therapy and research. By Robert B. Greenblatt, Robert B. Dienst, Herbert S. Kupperman, and Cecil R. Reinstein. September 1947. 8 pages. 5 cents.
290. Notes on the epidemiology of granuloma inguinale. By Charles Walter Clarke. September 1947. 6 pages. 5 cents.
291. "Penicillin-resistant gonorrhea" vs. "nonspecific urethritis." By George E. Parkhurst, Fred W. Harb, and George R. Cannefax. October 1947. 4 pages. 5 cents.
292. A study of preinoculation and preincubation factors in the primary isolation of *Neisseria gonorrhoeae*. By J. H. Schubert, Matthew A. Bucca, and J. D. Thayer. October 1947. 4 pages. 5 cents.
293. Two reports on out-patient attendance for treatment of syphilis, using penicillin in oil-beeswax. I. A study of clinic attendance. By Charles R. Hayman. II. Attendance record of patients treated by private physicians. By R. B. Aiken. October 1947. 4 pages. 5 cents.
294. A comparison of the efficiency of three common methods of transportation of gonorrheal specimens. By Lenore R. Peizer and Gustav I. Steffen. October 1947. 4 pages. 5 cents.
295. Penicillin therapy in early syphilis: III. By R. C. Arnold, J. F. Mahoney, John C. Cutler, and Sacha Levitan. November 1947. 4 pages. 5 cents.
296. Venereal disease control during the postwar period. By J. R. Heller, Jr. November 1947. 5 pages. 5 cents.
297. Quantitative serologic tests for syphilis. I. A standard method of reporting. By Ad Harris. November 1947. 3 pages. 5 cents.
298. Attempted immunization of rabbits against syphilis with killed *Treponema pallidum* and adjuvants. By Harold J. Magnuson, Seymour P. Halbert, and Barbara J. Rosenau. December 1947. 5 pages. 5 cents.
299. Contribution of the nurse in the schools to venereal-disease control. By Jane Barbara Taylor and Mildred F. Wills. December 1947. 5 pages. 5 cents.
300. Syphilis morbidity reporting by private physicians in the State of Florida. By R. F. Sondag and A. J. Sweeney. December 1947. 4 pages. 5 cents.

Supplement to the Journal of Venereal Disease Information

21. Autopsy studies in syphilis. A monograph. By Paul D. Rosahn. 1947. 67 pages. 25 cents.

Venereal Disease Bulletin

99. Venereal-disease control. A brief presentation of the venereal-disease control plan practiced in the United States of America. March 1947. 10 pages. 5 cents.

National Office of Vital Statistics Publications

Current mortality analysis (monthly), vol. 5, Nos. 5-9, 1947.

- ✓ A list of current publications of the National Office of Vital Statistics, August 1947, 6 pages.

Monthly marriage report (marriage licenses issued in major cities), new series: vol. 1, Nos. 6-11, 1947.

Monthly vital statistics bulletin, vol. 9, Nos. 5-10, 1947.

Quarterly marriage report (marriage licenses issued in the United States by State, 1947), vol. 2, Nos. 1-3.

Vital statistics—Special reports, vol. 25, national summaries:

No. 18. Plural birth statistics United States and each State, 1944. 337 to 362 pages.

No. 19. Index to volume 25. 363 to 370 pages.

Vital statistics—Special reports, vol. 27, national summaries:

No. 3. Deaths and death rates for selected causes: United States, each division and State, 1945. 27 to 46 pages.

No. 4. Infant mortality by race and by urban and rural areas: United States, each division and State, 1945. 47 to 54 pages.

No. 5. Births and deaths by specified race: United States, each division and State, 1945. 55 to 58 pages.

No. 6. Annual summary of motor-vehicle accident fatalities, 1945. 59 to 130 pages.

No. 7. Maternal mortality by cause and by race: United States and each State, 1945. 131 to 146 pages.

No. 8. Births and deaths by urban and rural areas: United States, each division and State, 1945. 147 to 152 pages.

No. 9. Births by person in attendance, by race and by urban and rural areas: United States, each division and State, 1945. 153 to 162 pages.

No. 10. Marriage and divorce statistics: United States, 1946. 163 to 176 pages.

No. 11. Deaths and death rates for selected causes, by age, race, and sex: United States, 1945. 177 to 214 pages.

No. 12. Infant mortality from selected causes, by age, race, and sex: United States, 1945. 215 to 232 pages.

Vital Statistics—Special reports, vol. 23, selected studies:

No. 16. Is family size increasing? 317 to 326 pages.

No. 17. Seasonal variation in the crude birth rate. 327 to 336 pages.

Vital statistics—Special reports, vol. 26, State summaries:

Nos. 16-54. Summary of vital statistics, 1945, for each State, Kansas through Wyoming (issued in alphabetic order); and the possessions of the United States: Alaska, Hawaii, Puerto Rico, and the Virgin Islands. 201 to 674 pages.

Weekly Mortality Index, vol. 18, Nos. 26-51.

✓ Where to write concerning marriage and divorce records. August 1947. 4 pages.

Summary of international vital statistics, 1937-1944 (1947). 299 pages. \$1.25.

Uniform definitions of motor vehicle accidents, 1947. 17 pages.

INCIDENCE OF DISEASE

No health department, State or local, can effectively prevent or control disease without knowledge of when, where, and under what conditions cases are occurring

UNITED STATES

REPORTS FROM STATES FOR WEEK ENDED MAY 8, 1948

Summary

A total of 107 cases of poliomyelitis was reported for the current week, as compared with 56 last week and a 5-year (1943-47) median of 26, reported for the corresponding week last year. Of the current total, Texas reported 55 (last week 25), California 14 (last week 3), North Carolina 8 (last week 7), and Indiana and Iowa 4 each. The total for the 7-week period since the average date of seasonal low incidence is 321, as compared with a 5-year median of 196 (reported last year), 126 in 1944 (the lowest number reported for a corresponding period of the past 5 years) and 219 in 1945 (the highest).

The incidence of measles continued above the 5-year median. A total of 28,343 cases was reported, as compared with 28,426 last week and a 5-year median of 26,032. Of the current total, 14,279 cases, or 50 percent occurred in the Middle Atlantic and East North Central areas (last year 3,037, or 37 percent of the 8,228 cases reported for the week). The total for the year to date is 334,940, as compared with 107,221 for the same period last year and a 5-year median of 340,866.

Of the total of 9 cases of Rocky Mountain spotted fever (last week 2, 5-year median 6), Maryland reported 3, Wyoming 2, and North Carolina, Oklahoma, Colorado, and Oregon, 1 each. The total to date is 22 (the same as the 5-year median), as compared with 21 for the same period last year.

One case of smallpox was reported, in Louisiana, and 1 case of anthrax, in Pennsylvania. Of the 7 cases of leprosy reported, 5 were in Florida and 1 each in New York and California. One case of psittacosis was reported in California.

A total of 9,266 deaths from all causes was recorded during the week in 93 large cities in the United States, as compared with 9,041 last week, 9,190 and 9,144, respectively, for the corresponding weeks of 1947 and 1946, and a 3-year (1945-47) median of 9,147. The total for the year to date is 188,841, as compared with 189,114 for the corresponding period last year. Infant deaths totaled 655, as compared with 679 last week, 769 and 619, respectively, for the corresponding week of 1947 and 1946, and a 3-year median of 619. The cumulative figure is 13,072, as compared with 15,064 for the corresponding period last year.

Telegraphic morbidity reports from State health officers for the week ended May 8, 1948, and comparison with corresponding week of 1947 and 5-year median

In these tables a zero indicates a definite report, while leaders imply that, although none was reported, cases may have occurred.

Division and State	Diphtheria			Influenza			Measles			Meningitis, meningococcus		
	Week ended—		Median 1943-47	Week ended—		Median 1943-47	Week ended—		Median 1943-47	Week ended—		Median 1943-47
	May 8, 1948	May 3, 1947		May 8, 1948	May 3, 1947		May 8, 1948	May 3, 1947		May 8, 1948	May 3, 1947	
NEW ENGLAND												
Maine.....	1	1	1	2	4	—	34	134	134	0	0	1
New Hampshire.....	0	0	0	—	—	—	20	4	23	0	0	0
Vermont.....	0	0	0	—	—	—	31	192	145	0	0	0
Massachusetts.....	6	9	8	—	—	—	1,369	461	952	3	1	5
Rhode Island.....	0	1	1	—	—	—	1	182	28	0	0	1
Connecticut.....	0	1	1	4	9	—	119	842	476	0	1	2
MIDDLE ATLANTIC												
New York.....	16	20	15	(1)	16	16	2,420	531	1,624	8	8	15
New Jersey.....	0	15	5	5	4	4	1,805	420	1,252	1	2	3
Pennsylvania.....	2	17	10	(2)	(2)	21	2,188	305	818	6	3	13
EAST NORTH CENTRAL												
Ohio.....	7	9	9	—	8	8	1,447	920	734	2	2	8
Indiana.....	20	5	4	—	4	4	1,335	111	261	0	3	3
Illinois.....	0	2	4	2	12	12	1,550	185	719	6	4	15
Michigan ¹	4	3	5	—	16	3	1,455	128	1,067	3	1	4
Wisconsin.....	1	1	1	13	24	38	2,079	437	1,854	2	1	1
WEST NORTH CENTRAL												
Minnesota.....	3	9	6	—	2	—	645	460	390	2	0	3
Iowa.....	0	1	2	—	—	—	255	127	249	0	1	1
Missouri.....	4	1	2	2	3	3	281	26	179	1	5	5
North Dakota.....	2	1	1	—	6	27	19	9	6	4	1	1
South Dakota.....	1	1	1	—	—	—	86	71	55	0	0	0
Nebraska.....	0	0	1	2	8	3	355	11	157	2	1	0
Kansas.....	4	4	4	1	2	—	52	13	514	0	2	2
SOUTH ATLANTIC												
Delaware.....	1	1	0	—	—	—	27	2	30	0	0	0
Maryland ¹	3	1	7	1	2	4	429	52	223	1	0	3
District of Columbia.....	0	0	0	—	—	—	132	25	77	0	0	0
Virginia.....	4	3	4	201	893	143	217	282	452	4	2	7
West Virginia.....	7	1	3	16	24	13	130	31	45	0	7	5
North Carolina.....	10	9	6	—	—	—	19	115	353	0	3	3
South Carolina.....	4	4	2	239	652	229	168	277	271	0	1	3
Georgia.....	3	0	2	3	41	2	39	107	107	1	1	1
Florida.....	5	4	3	3	78	3	135	60	87	1	1	3
EAST SOUTH CENTRAL												
Kentucky.....	5	6	2	1	6	6	228	9	153	0	4	4
Tennessee.....	2	5	4	18	83	26	183	104	196	1	2	4
Alabama.....	4	0	4	10	127	21	18	285	212	1	2	2
Mississippi ¹	3	3	3	2	45	—	25	23	—	1	0	4
WEST SOUTH CENTRAL												
Arkansas.....	2	2	2	106	152	21	145	72	122	1	2	2
Louisiana.....	2	4	4	1	9	5	8	45	84	2	3	3
Oklahoma.....	0	0	4	44	85	34	97	6	52	1	1	2
Texas.....	15	17	23	392	938	512	2,789	414	647	4	4	9
MOUNTAIN												
Montana.....	0	0	0	7	42	6	121	140	105	0	0	0
Idaho.....	0	0	0	19	21	2	143	1	58	0	0	0
Wyoming.....	0	1	1	1	1	—	78	15	91	0	0	0
Colorado.....	4	9	9	3	74	18	678	141	299	4	0	1
New Mexico.....	0	1	1	—	12	3	29	62	62	0	0	0
Arizona.....	5	0	1	28	85	42	428	61	61	0	0	0
Utah ¹	2	1	0	2	28	13	306	10	179	0	0	1
Nevada.....	0	0	0	—	—	—	4	—	7	0	0	0
PACIFIC												
Washington.....	0	0	7	—	11	—	720	47	307	1	3	3
Oregon.....	2	2	2	15	36	16	157	33	191	0	0	3
California.....	4	13	20	14	33	29	3,344	240	1,267	4	7	15
Total.....	158	188	190	1,157	3,586	1,432	28,343	8,228	26,032	67	79	158
18 weeks.....	3,475	4,821	4,821	130,335	290,376	182,740	334,940	107,221	340,866	1,453	1,596	4,167
Seasonal low week ⁴	(27th) July 5-11			(30th) July 26-Aug. 1			(35th) Aug. 30-Sept. 5			(37th) Sept. 13-19		
Total since low.....	9,833	12,387	13,549	173,893	323,351	323,351	369,886	130,108	378,879	2,235	2,568	6,619

¹ New York City only. ² Philadelphia only.

³ Period ended earlier than Saturday.

⁴ Dates between which the approximate low week ends. The specific date will vary from year to year.

*Correction (deducted from cumulative total): Diphtheria, South Carolina, week ended Apr. 24, 7 cases (instead of 13).

Telegraphic morbidity reports from State health officers for the week ended May 8, 1948, and comparison with corresponding week of 1947 and 5-year median—Con.

Division and State	Poliomyelitis			Scarlet fever			Smallpox			Typhoid and para typhoid fever		
	Week ended—		Med- ian 1943- 47	Week ended		Med- ian 1943- 47	Week ended—		Med- ian 1943- 47	Week ended—		Med- ian 1943- 47
	May 8, 1948	May 3, 1947		May 8, 1948	May 3, 1947		May 8, 1948	May 3, 1947		May 8, 1948 ¹	May 3, 1947	
NEW ENGLAND												
Maine.....	0	0	0	17	10	18	0	0	0	0	1	0
New Hampshire.....	0	0	0	3	8	11	0	0	0	0	0	0
Vermont.....	0	0	0	1	6	9	0	0	0	0	0	0
Massachusetts.....	0	0	0	216	92	309	0	0	0	2	4	2
Rhode Island.....	0	0	0	4	9	16	0	0	0	0	0	0
Connecticut.....	0	2	0	20	30	62	0	0	0	1	0	1
MIDDLE ATLANTIC												
New York.....	1	6	2	* 237	355	553	0	2	0	2	5	2
New Jersey.....	2	0	0	52	120	153	0	0	0	0	0	0
Pennsylvania.....	0	0	0	330	204	436	0	0	0	5	5	5
EAST NORTH CENTRAL												
Ohio.....	1	1	1	223	231	320	0	0	0	4	12	4
Indiana.....	4	0	0	54	77	78	0	1	3	3	2	2
Illinois.....	2	0	0	96	99	198	0	0	0	3	17	2
Michigan.....	0	1	0	159	114	176	0	0	0	0	3	1
Wisconsin.....	0	0	0	52	66	193	0	3	1	0	1	0
WEST NORTH CENTRAL												
Minnesota.....	1	0	0	34	44	70	0	0	0	0	1	0
Iowa.....	4	1	1	15	23	57	0	0	0	0	0	0
Missouri.....	0	0	0	* 30	17	53	0	0	0	1	3	1
North Dakota.....	0	2	0	4	4	11	0	0	0	0	0	0
South Dakota.....	0	0	0	2	4	13	0	0	0	0	0	0
Nebraska.....	0	0	0	23	22	27	0	0	0	0	1	0
Kansas.....	1	0	0	23	40	71	0	0	0	0	1	0
SOUTH ATLANTIC												
Delaware.....	0	0	0	4	4	4	0	0	0	1	0	0
Maryland.....	0	0	0	* 22	41	136	0	0	0	1	0	1
District of Columbia.....	0	0	0	9	11	22	0	0	0	0	0	0
Virginia.....	0	0	0	13	20	61	0	0	0	0	0	1
West Virginia.....	0	0	0	* 22	14	25	0	0	0	1	1	1
North Carolina.....	8	1	1	10	20	37	0	0	0	1	0	2
South Carolina.....	2	0	0	1	3	3	0	0	0	4	1	1
Georgia.....	1	0	0	6	6	9	0	0	0	2	0	2
Florida.....	3	2	3	4	3	5	0	0	0	2	0	2
EAST SOUTH CENTRAL												
Kentucky.....	0	0	0	21	24	32	0	2	0	0	3	1
Tennessee.....	0	0	0	20	23	41	0	0	0	2	0	1
Alabama.....	3	0	0	9	9	9	0	0	0	3	1	1
Mississippi.....	0	0	0	2	7	5	0	0	0	0	2	1
WEST SOUTH CENTRAL												
Arkansas.....	0	1	0	8	2	7	0	0	0	4	1	2
Louisiana.....	1	0	0	1	5	6	1	0	0	3	2	2
Oklahoma.....	1	0	0	8	4	12	0	0	0	0	0	0
Texas.....	* 55	2	2	16	15	48	0	0	0	7	6	6
MOUNTAIN												
Montana.....	1	0	0	5	3	17	0	0	0	0	0	0
Idaho.....	1	1	0	3	3	31	0	0	0	0	0	0
Wyoming.....	0	0	0	1	7	12	0	0	0	0	0	0
Colorado.....	0	0	0	37	45	48	0	0	0	0	0	0
New Mexico.....	1	1	0	0	9	9	0	1	0	0	0	1
Arizona.....	0	0	0	4	7	13	0	0	0	0	0	0
Utah.....	0	0	0	17	18	22	0	0	0	0	0	0
Nevada.....	0	0	0	0	1	1	0	0	0	0	0	0
PACIFIC												
Washington.....	0	0	0	42	22	37	0	0	0	0	2	0
Oregon.....	0	0	0	20	16	23	0	0	0	0	1	0
California.....	14	5	3	90	130	197	0	0	0	4	6	4
Total.....	107	26	26	1,990	2,047	3,859	1	9	15	56	82	54
18 weeks.....	669	808	616	41,020	47,007	71,761	44	111	191	855	820	1,029
Seasonal low week.....	(11th) Mar. 15-21			(32nd) Aug. 9-15			(35th) Aug. 30- Sept. 5			(11th) Mar. 15-21		
Total since low.....	321	196	196	63,559	73,693	110,082	65	165	272	382	335	425

¹ Period ended earlier than Saturday.

² Dates between which the approximate low week ends. The specific date will vary from year to year.

³ Including paratyphoid fever and salmonella infections reported separately, as follows: Massachusetts 2, Connecticut 1, New York 1, Pennsylvania 1, Indiana 1, West Virginia 1, South Carolina 1, Georgia 1, Tennessee 1, Texas 1.

⁴ Including cases reported as streptococcal sore throat.

⁵ 19 of these cases were for a 4-week period.

Telegraphic morbidity reports from State health officers for the week ended May 8, 1948, and comparison with corresponding week of 1947 and 5-year median—Con.

Division and State	Whooping cough			Week ended May 8, 1948								
	Week ended—		Me- dian 1943- 47	Dysentery			En- ceph- alitis, infec- tious	Rocky Mt. spot- ted fever	Tula- remia	Ty- phus fever, en- demic	Un- du- lant fever	
	May 8, 1948	May 3, 1947		Ame- bic	Bacil- lary	Un- spec- ified						
NEW ENGLAND												
Maine.....	11	31	36	—	—	—	—	—	—	—	—	
New Hampshire.....	—	3	3	—	—	—	—	—	—	—	—	
Vermont.....	21	7	12	—	—	—	—	—	—	—	—	
Massachusetts.....	39	104	135	—	2	—	2	—	—	—	—	
Rhode Island.....	4	24	18	—	—	—	—	—	—	—	—	
Connecticut.....	19	44	29	—	—	—	—	—	—	—	—	
MIDDLE ATLANTIC												
New York.....	95	174	174	11	—	—	—	—	—	2	5	
New Jersey.....	40	145	132	—	—	—	—	—	—	—	2	
Pennsylvania.....	101	158	158	—	—	—	—	—	—	—	1	
EAST NORTH CENTRAL												
Ohio.....	47	176	154	—	10	—	—	—	—	—	6	
Indiana.....	36	42	28	1	—	—	1	—	—	—	3	
Illinois.....	55	102	99	5	1	—	4	—	—	—	8	
Michigan ¹	40	260	132	7	—	—	—	—	—	—	4	
Wisconsin.....	96	153	85	—	—	—	—	—	—	—	12	
WEST NORTH CENTRAL												
Minnesota.....	12	32	12	—	—	—	—	—	—	—	1	
Iowa.....	8	24	20	—	—	—	—	—	—	—	2	
Missouri.....	13	22	22	—	—	—	—	—	—	—	—	
North Dakota.....	2	2	2	—	—	—	—	—	—	—	—	
South Dakota.....	1	—	—	—	—	—	—	—	—	—	—	
Nebraska.....	3	37	8	1	—	—	2	—	—	—	—	
Kansas.....	84	25	36	—	1	—	—	—	—	—	9	
SOUTH ATLANTIC												
Delaware.....	1	5	1	—	—	—	—	—	—	—	1	
Maryland ¹	16	80	73	—	—	4	1	3	—	—	2	
District of Columbia.....	5	9	9	—	—	—	—	—	—	—	—	
Virginia.....	33	85	63	1	—	21	—	—	—	—	1	
West Virginia.....	1	47	32	—	—	—	—	—	—	—	—	
North Carolina.....	75	66	97	—	—	—	—	1	—	—	—	
South Carolina.....	82	109	68	1	1	—	2	—	—	—	—	
Georgia.....	12	27	14	1	1	—	—	—	3	—	4	
Florida.....	14	77	42	4	—	5	—	—	—	2	1	
EAST SOUTH CENTRAL												
Kentucky.....	4	53	39	—	—	—	—	—	—	—	—	
Tennessee.....	26	36	33	2	—	1	—	—	2	—	1	
Alabama.....	22	64	37	—	—	—	—	—	—	2	2	
Mississippi ¹	3	6	—	2	—	—	—	—	—	—	2	
WEST SOUTH CENTRAL												
Arkansas.....	31	45	16	3	1	—	—	—	4	—	—	
Louisiana.....	1	9	9	3	—	—	—	—	2	1	1	
Oklahoma.....	38	20	20	—	—	3	—	1	1	—	1	
Texas.....	390	763	270	24	334	44	—	—	2	5	9	
MOUNTAIN												
Montana.....	6	5	5	—	—	—	—	—	—	—	1	
Idaho.....	5	9	9	—	—	—	—	—	—	—	—	
Wyoming.....	—	21	6	—	—	—	—	2	1	—	2	
Colorado.....	38	42	35	—	1	—	—	1	—	—	2	
New Mexico.....	12	28	9	—	—	—	—	—	—	—	—	
Arizona.....	30	34	27	—	—	30	—	—	—	—	—	
Utah ¹	19	5	44	—	—	—	—	—	—	—	—	
Nevada.....	1	—	—	—	—	—	—	—	—	—	—	
PACIFIC												
Washington.....	36	26	28	—	—	—	—	—	—	—	1	
Oregon.....	15	22	19	3	—	—	—	1	—	—	1	
California.....	69	351	313	5	8	—	2	—	1	1	5	
Total.....	1,712	3,609	2,646	84	350	108	14	9	16	13	93	
Same week, 1945.....	3,609			46	230	117	7	3	15	26	117	
Average, 1943-45.....	2,646			26	285	109	11	6	9	45	112	
52 weeks: 1948.....	38,450			1,258	5,107	3,205	158	22	324	247	1,640	
1947.....	48,000			828	5,303	3,619	121	21	570	690	1,868	
Average, 1943-47.....	44,726			530	5,261	1,868	153	22	294	817	1,638	

¹ Period ended earlier than Saturday.

² 3-year median 1945-47.

Anthrax: Pennsylvania 1 case. Leprosy: New York 1, Florida 5, California 1. Psittacosis: California 1 case.

Alaska, week ended May 1: Influenza 2, pneumonia 2, whooping cough 19; week ended May 8, influenza 4, pneumonia 4, whooping cough 18.

Territory of Hawaii, week ended May 8: Rabies 0, leprosy 1, measles 1, scarlet fever 1, whooping cough 14.

WEEKLY REPORTS FROM CITIES *

City reports for week ended May 1, 1948

This table lists the reports from 87 cities of more than 10,000 population distributed throughout the United States, and represents a cross section of the current urban incidence of the diseases included in the table.

Division, State, and city	Diphtheria cases	Encephalitis, infectious, cases	Influenza		Measles cases	Meningitis, meningococcus, cases	Pneumonia deaths	Poliomyelitis cases	Scarlet fever cases	Smallpox cases	Typhoid and paratyphoid fever cases	Whooping cough cases
			Cases	Deaths								
NEW ENGLAND												
Maine:												
Portland	0	0		0		0	3	0	0	0	0	7
New Hampshire:												
Concord	0	0		0		0	1	0	0	0	0	
Vermont:												
Barre	0	0		0		0	0	0	0	0	0	
Massachusetts:												
Boston	1	0		0	384	0	5	0	69	0	1	5
Fall River	0	0		0	25	0	0	0	2	0	0	5
Springfield	0	0		0	3	0	2	0	1	0	0	
Worcester	0	0		0	27	0	2	0	5	0	0	4
Rhode Island:												
Providence	0	0	2	0	4	0	3	0	3	0	1	15
Connecticut:												
Hartford	0	0		0	4	0	0	0	4	0	0	
New Haven	0	0	1	0	4	0	1	0	3	0	0	4
MIDDLE ATLANTIC												
New York:												
Buffalo	0	0		0	28	0	7	0	11	0	0	5
New York	6	2	1	1	1,586	2	68	1	82	0	2	29
Rochester	0	0		0	2	0	1	0	2	0	1	
Syracuse	0	0		0	3	0	0	0	1	0	0	10
New Jersey:												
Camden	0	0		0	27	0	2	0	1	0	0	
Newark	0	0	2	0	263	0	1	0	9	0	0	4
Trenton	0	0	1	0	2	0	3	0	7	0	0	
Pennsylvania:												
Philadelphia	2	0	1	1	873	3	23	0	43	0	0	7
Pittsburgh	0	0		0	2	3	10	0	70	0	0	3
Reading	0	0		0	3	0	2	0	15	0	0	1
EAST NORTH CENTRAL												
Ohio:												
Cincinnati	1	0		0	111	1	14	0	12	0	2	2
Cleveland	1	0		1	21	0	4	0	38	0	1	
Columbus	0	0		0	35	1	2	0	4	0	0	2
Indiana:												
Fort Wayne	0	0		0	14	0	0	0	1	0	0	
Indianapolis	0	0		0	164	1	9	0	11	0	0	5
South Bend	0	0		0	4	0	0	0	1	0	0	
Terre Haute	0	0		0		0	2	0	1	0	0	
Illinois:												
Chicago	1	0		2	690	1	25	0	37	0	2	13
Springfield	0	0		0	3	0	0	0	4	0	0	
Michigan:												
Detroit	1	1	1	0	519	1	6	0	67	0	0	16
Flint	0	0		0		1	1	0	5	0	0	
Grand Rapids	0	0		0	13	0	0	0	3	0	0	5
Wisconsin:												
Kenosha	0	0		0	84	0	0	0	1	0	0	
Milwaukee	0	0		0	111	0	3	0	24	0	0	3
Racine	0	0		0	53	0	0	0	3	0	0	5
Superior	0	0		0	227	0	0	0	0	0	0	1
WEST NORTH CENTRAL												
Minnesota:												
Duluth	0	0		0	284	0	1	0	1	0	0	
Minneapolis	1	0		0	49	0	5	0	9	0	0	4
St. Paul	0	0		0	61	2	3	0	4	0	0	10
Missouri:												
Kansas City	0	0	6	0	39	0	1	0	2	0	0	4
St. Joseph	0	0		0	6	1	0	0	0	0	0	5
St. Louis	2	0		0	203	0	6	0	7	0	0	6

* In some instances the figures include nonresident cases.

City reports for week ended May 1, 1948—Continued

Division, State, and city	Diphtheria cases	Encephalitis, infectious, cases	Influenza		Measles cases	Meningitis, meningococcus, cases	Pneumonia deaths	Pollomyelitis cases	Scarlet fever cases	Smallpox cases	Typhoid and paratyphoid fever cases	Whooping cough cases
			Cases	Deaths								
WEST NORTH CENTRAL—continued												
Nebraska:												
Omaha.....	0	0		0	175	0	3	0	0	0	0	
Kansas:												
Topeka.....	0	0		0	27	0	0	0	0	0	0	1
Wichita.....	0	0		0	5	0	4	0	1	0	0	27
SOUTH ATLANTIC												
Delaware:												
Wilmington.....	0	0		0	19	0	0	0	2	0	0	
Maryland:												
Baltimore.....	0	0	1	0	201	0	6	0	12	0	0	7
Cumberland.....	1	0		0		0	0	0	0	0	0	
Frederick.....	0	0		0		0	0	0	0	0	0	
District of Columbia:												
Washington.....	0	0		0	115	1	6	0	9	0	1	11
Virginia:												
Lynchburg.....	1	0		0	1	0	3	0	1	0	0	
Richmond.....	0	0		0	1	0	1	0	2	0	0	3
Roanoke.....	0	0		0		0	0	0	0	0	0	1
West Virginia:												
Charleston.....	0	0		0	2	0	4	0	0	0	0	
Wheeling.....	0	0		0	26	0	1	0	1	0	0	
North Carolina:												
Raleigh.....	0	0		0	1	0	0	0	0	0	0	2
Wilmington.....	0	0		0		0	0	0	2	0	0	7
Winston Salem.....	0	0		0	1	0	4	0	0	0	0	4
South Carolina:												
Charleston.....	0	0	13	0	2	0	1	0	1	0	0	2
Georgia:												
Brunswick.....	0	0		0		0	0	0	0	0	0	1
Savannah.....	0	0		0		0	0	0	0	0	0	1
Florida:												
Tampa.....	1	0		0	7	0	4	1	0	0	0	
EAST SOUTH CENTRAL												
Tennessee:												
Memphis.....	0	0		0	28	0	6	0	13	0	0	10
Nashville.....	0	0		0	5	0	1	0	3	0	0	
Alabama:												
Birmingham.....	3	0	1	1		6	1	1	1	0	0	7
Mobile.....	0	0		0		0	0	0	0	0	0	1
WEST SOUTH CENTRAL												
Arkansas:												
Little Rock.....	0	0		0	6	0	1	0	0	0	0	
Louisiana:												
New Orleans.....	2	0	1	0	7	0	5	0	1	0	3	1
Shreveport.....	0	0		0		0	3	0	0	0	0	
Oklahoma:												
Oklahoma City.....	0	0	4	0	22	0	3	0	2	0	0	3
Texas:												
Dallas.....	2	0		0	232	0	5	0	3	0	1	3
Galveston.....	0	0		0		0	2	1	0	0	0	
Houston.....	0	0	1	0	1	0	3	1	3	0	0	
San Antonio.....	1	0	1	0	37	0	3	0	1	0	0	
MOUNTAIN												
Montana:												
Billings.....	0	0		0	2	0	1	0	0	0	0	
Great Falls.....	0	0		0	2	0	0	0	0	0	0	
Helena.....	0	0		0		0	0	0	0	0	0	
Missoula.....	0	0		0		1	2	0	0	0	0	1
Colorado:												
Denver.....	0	0	1	0	288	0	1	1	4	0	0	13
Pueblo.....	0	0		0	9	0	0	0	1	0	0	1
Utah:												
Salt Lake City.....	1	0		0	71	0	0	0	0	0	0	1

City reports for week ended May 1, 1948—Continued

Division, State, and city	Diphtheria cases	Enecephalitis, infectious, cases	Influenza		Measles cases	Meningitis, meningococcus, cases	Pneumonia deaths	Poliomyelitis cases	Scarlet fever cases	Smallpox cases	Typhoid and paratyphoid fever cases	Whooping cough cases
			Cases	Deaths								
PACIFIC												
Washington:												
Seattle.....	0	0	-----	0	180	2	1	0	10	0	0	2
Spokane.....	0	0	-----	0	4	0	1	0	0	0	0	2
Tacoma.....	0	0	-----	0	25	0	0	0	2	0	0	-----
California:												
Los Angeles.....	1	0	3	0	331	0	7	2	19	0	2	12
Sacramento.....	1	0	-----	0	14	0	0	0	4	0	0	7
San Francisco.....	4	0	-----	0	221	1	8	0	10	0	0	7
Total.....	34	3	41	6	8,089	28	307	8	671	0	17	325
Corresponding week, 1947 ¹	68	-----	143	23	2,157	-----	390	-----	581	0	15	764
Average 1943-47.....	67	-----	84	19	6,400	-----	337	-----	1,435	1	12	660

¹ Exclusive of Oklahoma City.² 3-year average, 1945-47.³ 5-year median, 1943-47.

Rates (annual basis) per 100,000 population, by geographic groups, for the 87 cities in the preceding table (latest available estimated population, 84,123,200)

	Diphtheria case rates	Enecephalitis, infectious, case rates	Influenza		Measles case rates	Meningitis, meningococcus, case rates	Pneumonia death rates	Poliomyelitis case rates	Scarlet fever case rates	Smallpox case rates	Typhoid and paratyphoid fever case rates	Whooping cough case rates
			Case rates	Death rates								
New England.....	2.8	0.0	8.5	0.0	1,276	0.0	48.1	0.0	246	0.0	5.7	113
Middle Atlantic.....	3.7	0.9	2.3	2.3	1,391	3.7	54.2	0.5	112	0.0	1.4	27
East North Central.....	2.4	0.6	0.6	1.8	1,246	3.6	40.1	0.0	129	0.0	3.0	36
West North Central.....	6.0	0.0	12.1	0.0	1,707	6.0	46.3	0.0	48	0.0	0.0	115
South Atlantic.....	5.4	0.0	25.3	0.0	841	1.8	54.1	1.8	54	0.0	1.8	70
East South Central.....	17.7	0.0	5.9	5.9	195	35.4	47.2	5.9	100	0.0	0.0	106
West South Central.....	12.7	0.0	17.8	0.0	775	0.0	63.5	5.1	25	0.0	10.2	18
Mountain.....	8.3	0.0	8.3	0.0	3,073	8.3	33.0	2.3	41	0.0	0.0	132
Pacific.....	9.5	0.0	4.7	0.0	1,226	4.7	26.9	2.2	71	0.0	3.2	47
Total.....	5.2	0.5	6.3	0.9	1,239	4.3	47.0	1.2	103	0.0	2.6	50

Dysentery, amebic.—Cases: New York 5; Detroit 1; New Orleans 4; San Antonio 1; Los Angeles 3.

Dysentery, bacillary.—Cases: Worcester 1; New York 1; Chicago 2; Charleston, S. C. 1; Birmingham 1.

Dysentery, unspecified.—Cases: San Antonio 7.

PLAGUE INFECTION IN ARIZONA, NEW MEXICO, AND WASHINGTON

Under date of May 4, 1948, plague infection was reported proved in pools of fleas from rats and mice in Arizona, New Mexico, and Washington as follows:

ARIZONA

Navajo County.—A pool of 42 fleas from 4 wood rats, *Neotoma albigula*, trapped April 15 along State Highway No. 173 about 1½ miles southeast of Pinetop.

NEW MEXICO

Lincoln County.—A pool of 20 fleas from 21 wood rats, *Neotoma albigula*, trapped April 15, 1 mile north on a ranch road from a location 3 miles east of Capitan on U. S. Highway No. 380.

WASHINGTON

Kittitas County.—A pool of 180 fleas from 70 field mice, *Microtus* sp., trapped April 14 about 6 miles southeast of Kittitas.

TERRITORIES AND POSSESSIONS

Panama Canal Zone

Notifiable diseases—March 1948.—During the month of March 1948, certain notifiable diseases were reported in the Panama Canal Zone and terminal cities as follows:

Disease	Residence ¹									
	Panama City		Colon		Canal Zone		Outside the Zone and terminal cities		Total	
	Cases	Deaths	Cases	Deaths	Cases	Deaths	Cases	Deaths	Cases	Deaths
Chickenpox.....	4				2		8		14	
Diphtheria.....	8		1		3		5		17	
Dysentery:										
Amebic.....	4	1			2		6		12	1
Bacillary.....	1				4				5	
Hepatitis, infectious.....					1		3		4	
Malaria ²	3		3		3		113		122	
Measles.....	15		1		3		15		34	
Meningitis, meningococcus.....							1		1	1
Paratyphoid fever.....					1				1	
Pneumonia.....		11		1	14	1	3		14	16
Relapsing fever.....							1		1	
Streptococcal sore throat.....					1				1	
Tetanus.....							1	3	1	3
Tuberculosis.....		17		6	5	2	3		5	28
Typhoid fever.....							2	1	2	1

¹ If place of infection is known, cases are so listed instead of by residence.

² 8 recurrent cases.

³ In the Canal Zone only.

FOREIGN REPORTS

CANADA

Provinces—Communicable diseases—Week ended April 17, 1948.—During the week ended April 17, 1948, cases of certain communicable diseases were reported by the Dominion Bureau of Statistics of Canada as follows:

Disease	Prince Edward Island	Nova Scotia	New Brunswick	Que- bec	On- tario	Mani- toba	Sas- katch- ewan	Alber- ta	British Colum- bia	Total
Chickenpox.....		41	1	257	516	53	9	30	119	1,026
Diphtheria.....				11	2	1	1	2		17
German measles.....				36	21		2	8	12	79
Influenza.....		17			43	16			195	271
Measles.....		2		734	1,075	6	7	13	148	1,985
Meningitis, meningococcus.....					2					2
Mumps.....		19		373	290	73	107	49	41	952
Poliomyelitis.....					1		2			3
Scarlet fever.....		5	7	49	92		3	3		167
Tuberculosis (all forms).....		8	13	84	30	24	9	3	47	218
Typhoid and paratyphoid fever.....				5			1			6
Undulant fever.....					2			1		3
Venereal diseases:										
Gonorrhea.....	3	9	9	129	67	35	7	43	80	382
Syphilis.....	3	9	6	97	60	14	8	5	26	222
Other forms.....									2	2
Whooping cough.....				70	29	7	2	20	12	140

NORTHERN RHODESIA

Smallpox.—Information dated May 5, 1948, states that an epidemic of smallpox has been reported in the Zambesi Valley area, 100 miles east of Livingstone, Northern Rhodesia. The number of cases has not been ascertained, but 74 deaths are reported to have occurred during the past two months.

WORLD DISTRIBUTION OF CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER

From consular reports, international health organizations, medical officers of the Public Health Service, and other sources. The reports contained in the following tables must not be considered as complete or final as regards either the list of countries included or the figures for the particular countries for which reports are given.

CHOLERA

(Cases)

NOTE.—Since many of the figures in the following tables are from weekly reports, the accumulated totals are for approximate dates.

Place	January— February 1948	March 1948	April 1948—week ended—			
			3	10	17	24
AFRICA						
Egypt.....	1					
Cairo.....	1					
ASIA						
Burma.....	1	1				
India.....	18,243	6,778	1,848	1,288		
Ahmadabad.....	1	1				
Alleppey.....	1					
Calcutta ¹	1,326	854	282	343	292	435
Cawnpore.....		14	2	1	3	1
Cocanada.....	2					
Colachel.....		12				
Cuddalore.....	12					
Kilakarai.....	7	6				
Lucknow.....	6	3				
Madras.....	7	11	10	1		
Nagpur.....		1	1			
Negapatam.....	13	3				
New Delhi.....						1
Tuticorin.....	14	2				
India (French):						
Chandernagor.....	12	5				
Karikal.....	284					
Pondicherry.....	26	32				
Indochina (French):						
Cambodia.....	611	192		² 76		
Cochinchina.....	54	117		² 115	² 81	
Chaudoc.....	1					
Cholon.....	1	2		² 9		
Giadinh.....		4				
Longxuyen.....		7				
Mytho.....	4	8		² 8		
Rachgia.....	11	73		² 20		
Saigon.....	11	12	9	24	12	12
Laos.....	⁴ 12					
Tonkin.....		1				
Pakistan.....	5,441	609	1,289	1,433		
Chittagong.....	5	12				
Lahore.....		2			6	10
Siam.....	24	1				
Syria.....	3					

¹ Includes imported cases.

² For the period Apr. 1-10, 1948.

³ For the period Apr. 11-20, 1948.

⁴ Deaths.

PLAGUE

(Cases)

Place	January— February 1948	March 1948	April 1948—week ended—			
			3	10	17	24
AFRICA						
Belgian Congo.....	2	1	1			
British East Africa:						
Kenya.....	8	7				
Tanganyika.....	51	138				6
Madagascar.....	107	45		¹ 30		
Tananarive.....		10		¹ 6		
Rhodesia, Northern.....	13	8		1		
Union of South Africa.....	¹ 28	¹ 7		2		
ASIA						
Burma.....	¹ 296	106	18	15	9	6
Mandalay.....	5	1				
Rangoon.....	¹ 10	1			¹ 1	
China:						
Chekiang Province.....		3				
Wenchow.....		3				
Fukien Province.....	29	20				
Kiangsi Province.....	8					
Kwangtung Province.....	9					
Yunnan Province.....	31					
India.....	8,432	5,314	1,119	477		
Indochina (French):						
Annam.....	106	25		¹ 2		
Cochinchina.....	9	14		¹ 14		
Laos.....		2				
Java.....	4					
Siam.....	88	11	2			
EUROPE						
Portugal: Azores.....	7					
SOUTH AMERICA						
Argentina:						
Buenos Aires Province.....	5					
Ecuador.....	19					
Chimborazo Province.....	1					
Loja Province.....	1					
Peru:						
Huacho Department.....	1					
Lima Department.....	5					
OCEANIA						
Hawaii Territory: Plague-infected rats.....	5					

¹ For the period Apr. 1-10, 1948.² Includes 2 cases of pneumonic plague.³ Includes 1 imported case.⁴ Includes 2 imported cases.⁵ Imported.

SMALLPOX

(Cases)

(P = present)

AFRICA						
Algeria.....	72	38				
Angola.....	54					
Basutoland.....	3					
Belgian Congo.....	¹ 387	¹ 240				
British East Africa:						
Kenya.....	48	17				
Nyasaland.....	861	341	13	87		
Tanganyika.....	154	179	22	1		
Uganda.....	81	16	14			
Cameroon (French).....		2				
Dahomey.....	76	106		¹ 3	¹ 7	

See footnotes at end of table.

SMALLPOX—Continued

Place	January— February 1948	March 1948	April 1948—week ended—			
			3	10	17	24
AFRICA—continued						
Egypt.....	82	124	30	22		
Eritrea.....		5				
French Equatorial Africa.....	10					
French Guinea.....	48	20				
French West Africa: Haute-Volta.....	245	37				
Gambia.....	15	3			2	
Gold Coast.....	324	113	74			
Ivory Coast.....	164	35		1 20		
Libya.....	54	23	12	3	4	
Mauritania.....	1					
Morocco (French).....	11	2				
Mozambique.....	17	5				
Nigeria.....	316					
Niger Territory.....	147	73				
Rhodesia:						
Northern.....	103	3	1			
Southern.....	132					
Senegal.....	3					
Sierra Leone.....	74	9				
Sudan (Anglo-Egyptian).....	1 291	1 68	1 23	1 3	1 15	1 36
Sudan (French).....	8	6				
Swaziland.....			1			
Togo (British).....	9					
Togo (French).....	12	21				
Tunisia.....	399	66				
Union of South Africa.....	P	3				
ASIA						
Arabia.....	4 1	4 1				
British North Borneo.....	1					
Burma.....	682	628	148	135	104	156
Ceylon.....	4 6	4 3				
China.....	1, 490	677	95	115	97	69
India.....	15, 108	7, 686	1, 964	1, 919		
India (French).....	3	2				
Indochina (French).....	1, 024	548		1 139	1 41	
Iran.....	303	51				
Iraq.....	174	176	49	58	33	30
Japan.....	6	2		3		
Lebanon.....	57					
Malay States (Federated).....	264	49	6	2		
Manchuria.....	30					
Pakistan.....	4, 967	839	98			
Palestine.....	8					
Siam.....	304	87		3		
Straits Settlements.....						3
Syria.....	15	14		1		
Trans-Jordan.....		6				
EUROPE						
France.....		3				
Germany.....	1					
Portugal.....	26	25	2	3		
Spain.....	17					
Canary Islands.....	9					
NORTH AMERICA						
Guatemala.....	1					
Mexico.....	138	28				
SOUTH AMERICA						
Bolivia.....	16	15				
Brazil.....	9	1				
Chile.....		3			1	
Colombia.....	1, 018	879	32	28	31	
Ecuador.....	1 1, 083	1 215				
Paraguay.....	1 49	3				
Peru.....		20				
Venezuela.....	1 886	1 358		1 78	1 133	

1 Includes Alastrim.

2 For the period Apr. 1-10, 1948.

3 For the period Apr. 11-20, 1948.

4 Imported.

TYPHUS FEVER*

[Cases]

Place	January- February 1948	March 1948	April 1948—week ended—			
			3	10	17	24
AFRICA						
Algeria.....	54	28				
Basutoland.....	2	3	1			
Belgian Congo.....	43	16				
British East Africa:						
Kenya ¹	17	1	2	1		
Egypt.....	23	35	17	2		2
Eritrea.....	13	2	2			
Libya.....	46	28	11	11	20	
Morocco (French).....	22	21		3		
Nigeria ¹	2					
Senegal.....	3 1	3 1				
Sierra Leone.....	3 1	3 1				
Tunisia ¹	112	118				
Union of South Africa ¹	1	30				
ASIA						
Burma.....	5					
China ¹	33	3	1	1		
Indochina (French) ¹	3	3				
Iran ¹	41	10		3		
Iraq.....	32	19	2	14	3	6
Japan.....	182	55	7	17		
Manchuria.....	5					
Palestine ¹	12					
Philippine Islands.....	3 1					
Straits Settlements ¹	1	1			1	
Syria ¹	2	11			2	
Trans-Jordan.....	18	1			7	
Turkey (see Turkey in Europe).						
EUROPE						
Bulgaria.....	91	184	33			
Czechoslovakia.....	2	3	1			
France.....			3 1			
Germany.....	4					
Great Britain: Island of Malta ²	7	1				
Greece ¹	35	12	2	3		3
Hungary.....	22	26	2	1	1	
Italy ¹	9	11				
Luxemburg ⁴						
Netherlands.....	3 1					
Poland.....	62	21				
Rumania ¹	16,745	1,952				
Spain.....		1				
Turkey.....	117	35	15	8	11	5
Yugoslavia.....	107	151				
NORTH AMERICA						
Costa Rica ³	1					
Cuba ³	4	6	1			
Guatemala.....	18					
Jamaica ³	2			1		
Mexico ¹	223	31	2	2		
Panama Canal Zone.....	1					
Puerto Rico ³	5	3				2
SOUTH AMERICA						
Bolivia.....		3 11	12	3		
Brazil.....	56	6			7	2
Chile ¹	10	6 66				
Colombia.....	561	264				
Curacao ³	8	3				
Ecuador ¹	92	17			7 7	
Venezuela.....	13	2		4		
OCEANIA						
Australia ³	26	17	5	3	3	

* Reports from some areas are probably murine type, while others probably include both murine and louse-borne types.

¹ Includes murine type.

² For the period Apr. 1-10, 1948.

³ Murine type.

⁴ The previous report of cases of typhus fever in Luxemburg was an error. The cases were later stated to have been typhoid fever.

⁵ Deaths.

⁶ For the period Jan. 1-Mar. 15, 1948.

⁷ For the period Apr. 1-15, 1948.

YELLOW FEVER

[C indicates cases; D, deaths]

Place	January— February 1948	March 1948	April 1948—week ended—			
			3	10	17	24
AFRICA						
Ivory Coast:						
Gagnoa.....	C	1				
Nigeria: Lagos Island ¹						
SOUTH AMERICA						
Colombia:						
Antioquia Department.....	C	25				
Boyaca Department.....	D	1				
Caldas Department.....	D	2				
Cundinamarca Department.....	D	7				
Intendencia of Meta.....	D	3				

¹ The case of yellow fever in Igbo Village, Lagos Island, Nigeria, reported on p. 592 of the Public Health Reports for Apr. 30, 1948, in the column for week ended March 6, has since been reported not confirmed.

² Includes deaths used as cases.

DEATHS DURING WEEK ENDED MAY 1, 1948

[From the Weekly Mortality Index, issued by the National Office of Vital Statistics]

	Week ended May 1, 1948	Correspond- ing week, 1947
Data for 93 large cities of the United States:		
Total deaths.....	9,041	8,977
Median for 3 prior years.....	8,974	
Total deaths, first 18 weeks of year.....	179,575	179,924
Deaths under 1 year of age.....	679	747
Median for 3 prior years.....	645	
Deaths under 1 year of age, first 18 weeks of year.....	12,417	14,295
Data from industrial insurance companies:		
Policies in force.....	71,068,300	67,286,612
Number of death claims.....	12,406	13,724
Death claims per 1,000 policies in force, annual rate.....	9.1	10.6
Death claims per 1,000 policies, first 18 weeks of year, annual rate.....	10.2	10.0

a
P
to
2

d
o
in
t
r

a
a
o
a

r
S
c

I
v

F

The PUBLIC HEALTH REPORTS, first published in 1878 under authority of an act of Congress of April 29 of that year, is issued weekly by the United States Public Health Service through the Division of Public Health Methods, pursuant to the following authority of law: United States Code, title 42, sections 241, 245, 247; title 44, section 220.

It contains (1) current information regarding the incidence and geographic distribution of communicable diseases in the United States, insofar as data are obtainable, and of cholera, plague, smallpox, typhus fever, yellow fever, and other important communicable diseases throughout the world; (2) articles relating to the cause, prevention, and control of disease; (3) other pertinent information regarding sanitation and the conservation of the public health.

The PUBLIC HEALTH REPORTS is published primarily for distribution, in accordance with the law, to health officers, members of boards or departments of health, and other persons directly or indirectly engaged in public health work. Articles of special interest are issued as reprints or as supplements, in which forms they are made available for more economical and general distribution.

Requests for and communications regarding the PUBLIC HEALTH REPORTS, reprints, or supplements should be addressed to the Surgeon General, United States Public Health Service, Washington 25, D. C. Subscribers should remit direct to the Superintendent of Documents, Washington 25, D. C.

Librarians and others should preserve their copies for binding, as the Public Health Service is unable to supply the general demand for bound copies. Indexes will be supplied upon request.

UNITED STATES GOVERNMENT PRINTING OFFICE, WASHINGTON, D. C. : 1948

For sale by the Superintendent of Documents, United States Government Printing Office, Washington 25, D. C.

Price 10 cents. Subscription price \$4.00 a year